

This is a repository copy of *Plasticity, and Its Limits, in Adult Human Primary Visual Cortex*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/96225/>

Version: Submitted Version

Article:

Haak, Koen V, Morland, Antony B orcid.org/0000-0002-6754-5545 and Engel, Stephen A (2015) Plasticity, and Its Limits, in Adult Human Primary Visual Cortex. Multisensory Research. 297–307. ISSN 2213-4808

<https://doi.org/10.1163/22134808-00002496>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Multisensory Research

Plasticity, and its Limits, in Adult Human Primary Visual Cortex

--Manuscript Draft--

Manuscript Number:		
Full Title:	Plasticity, and its Limits, in Adult Human Primary Visual Cortex	
Short Title:	Limits of plasticity in Adult Human Primary Visual Cortex	
Article Type:	Special Issue: Multimodality of Early Sensory Processing	
Corresponding Author:	Koen V. Haak Radboud Universiteit Nijmegen Nijmegen, NETHERLANDS	
Corresponding Author's Institution:	Radboud Universiteit Nijmegen	
First Author:	Koen V. Haak	
Order of Authors:	Koen V. Haak	
	Antony B. Morland	
	Stephen A. Engel	
Abstract:	<p>There is an ongoing debate about whether adult human primary visual cortex (V1) is capable of large-scale cortical reorganization in response to bilateral retinal lesions. Animal models suggest that the visual neural circuitry maintains some plasticity through adulthood, and there are also a few human imaging studies in support this notion. However, the interpretation of these data has been brought into question, because there are factors besides cortical reorganization that could also explain the results. Still, how reasonable would it be to accept that adult human V1 does not reorganize itself in the face of disease? Here, we discuss new evidence for the hypothesis that adult human V1 is not as capable of reorganization as in animals and juveniles, because in adult humans, cortical reorganization would come with costs that outweigh its benefits. These costs are likely functional and visible in recent experiments on adaptation—a rapid, short-term form of neural plasticity—where they prevent reorganization from being sustained over the long-term.</p>	
Keywords:	Neuroplasticity; Primary Visual Cortex; Human Adults; Costs	
Suggested Reviewers:		
Opposed Reviewers:		
Funding Information:	Medical Research Council (G0401339)	Antony B. Morland
	National Science Foundation (BCS1028584)	Stephen A. Engel

1
2
3
4 **Plasticity, and its Limits, in Adult Human Primary Visual Cortex**
5
6

7 **Koen V. Haak^{1,*}, Antony B. Morland² and Stephen A. Engel³**
8
9

10 ¹ Donders Institute for Brain, Cognition and Behaviour, Centre for Cognitive Neuroimaging,
11 Radboud University, Nijmegen, The Netherlands
12
13

14 ² Hull-York Medical School and Department of Psychology, University of York, York, United
15 Kingdom
16
17

18 ³ Department of Psychology, University of Minnesota, Minneapolis, United States
19
20
21
22
23

24
25
26
27 **Abstract**
28
29

30 There is an ongoing debate about whether adult human primary visual cortex (V1) is capable of
31 large-scale cortical reorganization in response to bilateral retinal lesions. Animal models suggest
32 that the visual neural circuitry maintains some plasticity through adulthood, and there are also a
33 few human imaging studies in support this notion. However, the interpretation of these data has
34 been brought into question, because there are factors besides cortical reorganization that could
35 also explain the results. Still, how reasonable would it be to accept that adult human V1 does not
36 reorganize itself in the face of disease? Here, we discuss new evidence for the hypothesis that
37 adult human V1 is not as capable of reorganization as in animals and juveniles, because in adult
38 humans, cortical reorganization would come with costs that outweigh its benefits. These costs
39 are likely functional and visible in recent experiments on adaptation—a rapid, short-term form of
40 neural plasticity—where they prevent reorganization from being sustained over the long-term.
41
42
43
44
45
46
47
48
49
50

51 **Keywords**
52
53

54 Neuroplasticity, Primary Visual Cortex, Human Adults, Costs
55
56
57
58
59

60 * correspondence should be addressed to: k.haak@donders.ru.nl
61
62
63
64
65

1
2
3
4 If the retina is damaged in both eyes, primary visual cortex (V1) no longer receives
5 input. It is widely hypothesized that neurons within the V1 cortical lesion
6 projection zone (LPZ) respond to the absence of incoming information by shifting
7 their receptive fields from the blind toward intact portions of the visual field (as
8 suggested by e.g., Kaas et al., 1990; Gilbert and Wiesel, 1992; Darian-Smith and
9 Gilbert, 1994; Chino et al., 1992; Chino et al., 1995; Calford et al., 1999; Calford
10 et al., 2000; Schmid et al., 1995; Darian-Smith and Gilbert, 1995). This type of
11 visual brain plasticity is often called ‘remapping’, because the relocation of
12 neuronal receptive fields will change V1’s retinotopic map: retinotopic maps in
13 cortex emerge from the fact that nearby visual neurons have receptive fields at
14 nearby locations in the visual field, so when neurons shift the location of their
15 receptive fields, the retinotopic map will change accordingly. Visual brain
16 plasticity in the form of remapping is generally thought to be slow, because it
17 would require that structural changes are made to the underlying neural circuitry.
18 However, recently, the interpretation of the data presented in support of cortical
19 remapping has been seriously questioned (see Wandell and Smirnakis, 2009 for a
20 review).

21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42 Another important type of visual brain plasticity is neuronal adaptation. In contrast
43 to cortical remapping, neuronal adaptation is generally very fast (e.g., Clifford et
44 al., 2000; Kohn, 2007; Wandell and Smirnakis, 2009). For instance, if a person
45 looks at a waterfall for a short period of time and then shifts her gaze away, the
46 world will perceptually move upward (Anstis et al., 1998). This motion after-effect
47 is thought to be due to the fact that motion-processing visual neurons adjust their
48 responsiveness, which likely reflects an attempt to optimize visual processing by
49 maintaining a state of equal time-averaged activity and decorrelated neuronal
50 stimulus sensitivity (Attneave, 1954; Andrews, 1964, Ullman and Schechtman,
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 1982; Barlow and Földiák, 1989; Anstis et al., 1998; Wainwright, 1999; Benucci
5 et al., 2013; Haak et al., 2014a). Unlike the neuronal receptive field changes
6 underlying cortical remapping, the changes underlying neural adaptation do not
7 appear to rely on structural changes in the visual neural circuitry (Clifford et al.,
8 2000; Kohn, 2007; Wandell and Smirnakis, 2009), but rather on functional
9 interactions, such as mutual inhibition (Movshon and Lennie, 1979; Barlow and
10 Földiák, 1989).

11
12
13
14
15
16
17
18
19
20 Whereas neuronal adaptation is a firmly established property of the visual system
21 across species and ages (Clifford et al., 2000; Kohn, 2007), cortical remapping
22 following retinal lesions has yet to receive such status, particularly in adult humans
23 (for a review, see Wandell and Smirnakis, 2009). A handful of functional magnetic
24 resonance imaging (fMRI) studies have characterized abnormal functional
25 magnetic resonance imaging (fMRI) activity in V1 following retinal lesions (e.g.,
26 Baker et al., 2005; Schumacher et al., 2008; Baker et al., 2008; Dilks et al., 2009;
27 Dilks et al., 2014), thereby claiming evidence of large-scale reorganization.
28 However, abnormal activity in itself does not warrant the conclusion that cortical
29 remapping has occurred (Masuda et al., 2008; Wandell and Smirnakis, 2009;
30 Baseler et al., 2009; Masuda et al., 2010; Haak et al., 2014c). Cortical remapping
31 can only be concluded on the basis of abnormal activity patterns if it can also be
32 shown that the absence of visual input in itself does not change the measurements
33 in the same way. For instance, in a recent study, Baseler et al. (2011) compared the
34 fMRI activity patterns from the cortical lesion projection zone (LPZ) in macular
35 degeneration patients with real retinal lesions and healthy controls with simulated
36 retinal lesions. They found that although the LPZ in V1 was largely silenced by
37 both real and simulated retinal lesions, for a small fraction of voxels (~5%) the
38 patients with real retinal lesions exhibited the same apparent neuronal receptive
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 field changes as controls with simulated lesions, indicating that these changes were
5 caused by the absence of visual stimulation alone. Importantly, these voxels were
6 found far into the LPZ, indicating that the receptive field changes could not be
7 easily explained by measurement artifacts at the fringe of the LPZ (Haak et al.,
8 2012; Binda et al., 2013). Rather, they appeared to be a feature of visual cortical
9 processing, unveiled in the absence of visual stimulation.

10
11 In the first instance, it seems quite reasonable to expect that the adult human brain
12 would adjust itself in the face of retinal lesions. After all, if children are capable of
13 developing relatively normal vision even when an entire occipital lobe failed to
14 develop (e.g., Werth, 2006, Muckli et al., 2009, but see Haak et al., 2014c), one
15 might also expect the brain to at least ameliorate the consequences of a retinal
16 lesion by dedicating the now-redundant resources of deafferented cortex to
17 processing retinal inputs that are still intact. The net effect of such cortical
18 remapping would be quite similar to the perceptual ‘filling-in’ of the blind-spot of
19 the healthy retina, as well as the perceptual filling-in that occurs when someone
20 stares steadily at an image with patches of missing ‘texture’ for a prolonged period
21 of time (Ramachandran and Gregory, 1991; Pettet and Gilbert, 1992; Komatsu,
22 2006; Weil and Rees, 2011). Here, the patches of missing ‘image data’ will be
23 perceptually filled-in with the texture from surrounding image regions. Just like the
24 after-effects of neuronal adaptation, perceptual filling-in occurs fairly rapidly, and
25 the neural mechanisms underlying perceptual filling-in could likewise be
26 interpreted as a way of optimizing visual processing in the context of the preceding
27 stimulus history (i.e., in the vein of Horace Barlow’s redundancy reduction
28 hypothesis; c.f. Barlow, 1961; Barlow, 2001). Thus, perceptual filling-in, which
29 would be the consequence of cortical remapping in response to retinal lesions if it
30 occurred, seems to share important features with visual neuroplasticity in the form

1
2
3
4 of neuronal adaptation. Therefore, as the loss of vision endures, it might be
5 expected that the brain would eventually change its neural circuitry via cortical
6 remapping to sustain perceptual filling-in over the long-term. However, this does
7 not appear to be the case—perceptual filling-in does not appear to produce long-
8 term, hard-wired changes in the face of a retinal lesion.
9

10
11 Interestingly, recent work now suggests that neuronal adaptation may also not
12 produce hard-wired changes over the long-term. Using immersive virtual reality,
13 Haak et al. (2014b) exposed a group of young adults to a world with only very
14 little vertical visual contrast energy for four days continuously, in an attempt to
15 mimic classic selective rearing experiments (Hirsch and Spinelli, 1970; Blakemore
16 and Cooper, 1970) in adult humans. Just as staring at a waterfall for a prolonged
17 period of time changes the response gains of motion-sensitive neurons, the
18 prolonged viewing of a world with relatively little vertical contrast will cause
19 adjustments to the responsiveness of orientation-selective cells in primary visual
20 cortex (e.g., Graham, 1989; Maffei et al., 1973; Movshon and Lennie, 1973;
21 Ohzawa et al., 1985; Dragoi et al., 2000). Monitoring for the perceptual
22 consequences of these changes in the responsiveness of orientation-selective
23 neurons, Haak et al. (2014b) found that adaptation increased in magnitude during
24 the first day, but then decreased, despite the sustained presence of the adapting
25 environment. Thus, it appears that there are factors that prevent visual
26 neuroplasticity, in the form of neuronal adaptation, from being sustained over the
27 long-term.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52

53 Haak et al. (2014b) concluded that if neuronal adaptation does in fact optimize
54 vision, then the decline in adaptation strength must be due to costs that outweighed
55 its benefits. An obvious candidate cost is the ‘coding catastrophe’, where changes
56 in the firing of neurons responsible for early visual processing are mistaken for
57
58
59
60
61
62
63
64
65

1
2
3
4 stimulus-changes by neurons that are responsible for subsequent, higher-level
5 stages of visual processing (Schwartz et al., 2007; Series et al., 2009; Druv and
6 Carandini, 2014; Patterson et al., 2014). Indeed, when Patterson et al. (2014)
7 induced adaptive changes in the responsiveness of V1 neurons in the macaque by
8 having them stare at a drifting grating pattern for some time, this impeded the
9 ability of neurons in area MT, which receives most of its inputs from V1, to
10 integrate the V1 signals into a plaid pattern. They concluded that “the effects of
11 adaptation cascade through the visual system, derailing the downstream
12 representation of distinct stimulus attributes”.

13
14
15
16
17
18
19
20
21
22
23
24 Importantly, the decline in adaptation strength observed by Haak et al. (2014b) was
25 then followed by an increase in adaptation during subsequent days, indicating that
26 a second, more slowly acting adaptive mechanism was able to overcome the costs
27 of the initial adjustments in neuronal responsiveness. It is likely that this second,
28 slower form of adaptation reflects a process more similar to ‘perceptual learning’,
29 during which the visual system typically adjusts the neural codes in later rather
30 than in earlier visual areas (see e.g., Hochstein and Ahissar, 2002; Ahissar and
31 Hochstein, 2004). This would make sense, because later stages of visual processing
32 are typically concerned with more abstract sensory representations that require
33 more flexible neural codes. Moreover, compared with the areas responsible for the
34 early stages of visual processing, there are far fewer downstream areas that depend
35 on the information throughput of later visual areas. Thus, by shifting the adaptive
36 neural code adjustments from the early to higher-level visual areas, the brain may
37 be able to optimize vision without the adverse effects of the coding catastrophe.
38 Indeed, Haak et al. (2014b) observed that the tilt-aftereffect, an illusion thought to
39 be due to the coding-catastrophe, began to decline toward the end of the
40 experiment.

Here, we put forward the hypothesis that the same principles may also apply to visual processing in the face of retinal lesions. That is, the retinal lesion would initially cause rapid ‘adaptive’ changes in the location of the receptive fields of the neurons within the lesion projection zone in V1. These changes, however, will likely cause havoc at later stages of vision, due to the mismatch between the assumed and true stimulus locations that their afferents encode. Thus, early stage changes could soon be undone, as perceptual filling-in is postponed to later stages of visual processing (in line with e.g. De Weerd et al., 1995; Murakami et al., 1997; Cornelissen et al., 2006). A cost-induced shift of cortical reorganization from earlier to later stages of visual processing would straightforwardly explain why reorganization in the form of cortical remapping appears to be absent in adult V1 (Sunness et al., 2004; Smirnakis et al., 2005; Wandell and Smirnakis, 2009; Baseler et al., 2011). The hypothesis is also consistent with the task-dependent differences in V1 activation between patients and controls reported by Masuda et al. (2008, 2010), who reasoned that the task-dependent responses in the LPZ of V1 must reflect unmasked feedback signals from the extra-striate visual areas, rather than cortical reorganization at the level of V1.

As there are many controversies surrounding the claims of cortical remapping in cats, primates and humans (see Wandell and Smirnakis, 2009), perhaps the most convincing evidence of cortical remapping in adult V1 comes from studies in mice (e.g., Keck et al., 2008). Why would the coding catastrophe not limit cortical remapping in adult mice with binocular retinal lesions? One reason could be that mice do not rely as much on vision as cats, primates and humans do (mice are nocturnal and navigate using mainly their nose and whiskers), and so the costs of cortical remapping may be lower for them. Another reason could be that V1 is in fact a relatively high-level area within the mouse visual hierarchy, such that it

1
2
3
4 serves relatively late-stage rather than early-stage visual processing. If so, fewer
5 visual processing stages would be adversely affected by plastic changes in V1. In a
6 similar vein, the costs of cortical remapping may be lower in mice if mouse V1
7 were to feed its information straight to the areas with highly flexible processing
8 capabilities (see e.g., Wang and Burkhalter, 2007), thereby bypassing mid-level
9 processing stages (corresponding to e.g. human visual area V2) that are more
10 constrained in the variety of their inputs. Regardless, there is no doubt that visual
11 cortex in mice is very different from visual cortex in (human) primates, with
12 mouse visual cortex containing for instance no orientation columns, and many
13 fewer extrastriate visual areas (Baker, 2013; Huberman and Niell, 2011), leaving
14 open the possibility that it contains different mechanisms of plasticity.
15
16
17
18
19
20
21
22
23
24
25
26
27

28 Claims of cortical remapping have not been limited to visual cortex. In auditory
29 cortex, for instance, ‘filling-in’ like changes have been observed in the topographic
30 cortical representation of sound frequency (i.e., auditory cortex’ tonotopic map)
31 starting weeks after localized lesions were applied to the cochlea of the inner ear
32 (e.g., Robertson and Irvine, 1989). Though the criticisms of cortical remapping in
33 the visual domain may also apply to auditory cortex, it is interesting to note that
34 primary auditory cortex (A1) is a relatively later stage within the auditory
35 processing pathways than V1 is within the visual processing hierarchy. That is,
36 there are many more subcortical stops before auditory information reaches cortex
37 than there are stops leading up to V1, and there are many fewer high-level cortical
38 auditory areas than there are high-level visual areas beyond V1. Indeed, it has been
39 proposed that A1 represents stimuli in a highly task-dependent fashion, thereby
40 affording a relatively high amount of learning-induced plasticity (e.g., Ohl and
41 Scheich, 2005; Polley et al., 2006). Thus, the difference in the degree of cortical
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 reorganization in A1 and V1 appears to fit well with the idea that the brain prefers
5 to make plastic changes at the later stages of sensory processing.
6
7

8
9 In conclusion, there appears to be converging evidence to suggest that adult human
10 primary visual cortex is not very susceptible to cortical remapping. Here, we have
11 put forward the hypothesis that this could be due to the costs associated with
12 making changes at the very root of the visual processing hierarchy. It would make
13 more sense for the visual system to make long-term structural adjustments at later
14 stages of visual processing, because the later stages have fewer dependencies that
15 may be adversely affected (Hochstein and Ahissar, 2002; Ahissar and Hochstein,
16 2004). This principle may not only apply to brain plasticity in the form of cortical
17 remapping in response to bilateral retinal lesions, but also when it comes to various
18 multimodal forms of cortical reorganization.
19
20
21
22
23
24
25
26
27
28
29
30

31 *Acknowledgments*

32

33
34 This work was supported by National Science Foundation Grant BCS1028584 and
35 the Medical Research Council (G0401339).
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

- Ahissar, M., & Hochstein, S. (2004). The reverse hierarchy theory of visual perceptual learning. *Trends Cogn Sci*, 8, 457-464.
- Andrews, D. P. (1964). Error-Correcting Perceptual Mechanisms. *Quarterly Journal of Experimental Psychology*, 16, 104-115.
- Anstis, S., Verstraten, F. A., & Mather, G. (1998). The motion aftereffect. *Trends Cogn Sci*, 2, 111-117.
- Attneave, F. (1954). Some informational aspects of visual perception. *Psychol Rev*, 61, 183-193.
- Baker, C. I., Dilks, D. D., Peli, E., & Kanwisher, N. (2008). Reorganization of visual processing in macular degeneration: replication and clues about the role of foveal loss. *Vision Res*, 48, 1910-1919.
- Baker, C. I., Peli, E., Knouf, N., & Kanwisher, N. G. (2005). Reorganization of visual processing in macular degeneration. *J Neurosci*, 25, 614-618.
- Baker, M. (2013). Neuroscience: Through the eyes of a mouse. *Nature*, 502, 156-158.
- Barlow, H. (2001). Redundancy reduction revisited. *Network*, 12, 241-253.
- Barlow, H. B. (1961). The coding of sensory messages. In W. H. Thorpe & O. L. Zangwill (Eds.), *Current problems in animal behaviour* (pp. 331-3360). Cambridge: Cambridge University Press.
- Barlow, H. B., & Földiák, P. (1989). Adaptation and decorrelation in the cortex. In R. Durbin, C. Miall & G. Mitchison (Eds.), *The computing neuron* (pp. 54-72). Workingham, England: Addison-Wesley.
- Baseler, H. A., Gouws, A., Haak, K. V., Racey, C., Crossland, M. D., Tufail, A., Rubin, G. S., Cornelissen, F. W., & Morland, A. B. (2011). Large-scale remapping of visual cortex is absent in adult humans with macular degeneration. *Nat Neurosci*, 14, 649-655.
- Baseler, H. A., Gouws, A., & Morland, A. B. (2009). The Organization of the Visual Cortex in Patients with Scotomata Resulting from Lesions of the Central Retina. *Neuro-Ophthalmology*, 33, 149-157.
- Benucci, A., Saleem, A. B., & Carandini, M. (2013). Adaptation maintains population homeostasis in primary visual cortex. *Nat Neurosci*, 16, 724-729.
- Binda, P., Thomas, J. M., Boynton, G. M., & Fine, I. (2013). Minimizing biases in estimating the reorganization of human visual areas with BOLD retinotopic mapping. *J Vis*, 13, 13.

- 1
2
3
4 Blakemore, C., & Cooper, G. F. (1970). Development of the brain depends on the visual
5 environment. *Nature*, 228, 477-478.
6
7
8 Calford, M. B., Schmid, L. M., & Rosa, M. G. (1999). Monocular focal retinal lesions induce
9 short-term topographic plasticity in adult cat visual cortex. *Proc Biol Sci*, 266, 499-507.
10
11 Calford, M. B., Wang, C., Taglianetti, V., Waleszczyk, W. J., Burke, W., & Dreher, B. (2000).
12 Plasticity in adult cat visual cortex (area 17) following circumscribed monocular lesions of
13 all retinal layers. *J Physiol*, 524 Pt 2, 587-602.
14
15
16
17 Chino, Y. M., Kaas, J. H., Smith, E. L., 3rd, Langston, A. L., & Cheng, H. (1992). Rapid
18 reorganization of cortical maps in adult cats following restricted deafferentation in retina.
19 *Vision Res*, 32, 789-796.
20
21
22 Chino, Y. M., Smith, E. L., 3rd, Kaas, J. H., Sasaki, Y., & Cheng, H. (1995). Receptive-field
23 properties of deafferentated visual cortical neurons after topographic map reorganization in
24 adult cats. *J Neurosci*, 15, 2417-2433.
25
26
27
28 Clifford, G. M., & Farmer, R. D. (2000). Medical therapy for benign prostatic hyperplasia: a
29 review of the literature. *Eur Urol*, 38, 2-19.
30
31
32 Cornelissen, F. W., Wade, A. R., Vladusich, T., Dougherty, R. F., & Wandell, B. A. (2006). No
33 functional magnetic resonance imaging evidence for brightness and color filling-in in early
34 human visual cortex. *J Neurosci*, 26, 3634-3641.
35
36
37 Darian-Smith, C., & Gilbert, C. D. (1994). Axonal sprouting accompanies functional
38 reorganization in adult cat striate cortex. *Nature*, 368, 737-740.
39
40
41 Darian-Smith, C., & Gilbert, C. D. (1995). Topographic reorganization in the striate cortex of the
42 adult cat and monkey is cortically mediated. *J Neurosci*, 15, 1631-1647.
43
44
45 De Weerd, P., Gattass, R., Desimone, R., & Ungerleider, L. G. (1995). Responses of cells in
46 monkey visual cortex during perceptual filling-in of an artificial scotoma. *Nature*, 377, 731-
47 734.
48
49
50 Dhruv, N. T., & Carandini, M. (2014). Cascaded effects of spatial adaptation in the early visual
51 system. *Neuron*, 81, 529-535.
52
53
54 Dilks, D. D., Baker, C. I., Peli, E., & Kanwisher, N. (2009). Reorganization of visual processing
55 in macular degeneration is not specific to the "preferred retinal locus". *J Neurosci*, 29, 2768-
56 2773.
57
58
59
60
61
62
63
64
65

- Dilks, D. D., Julian, J. B., Peli, E., & Kanwisher, N. (2014). Reorganization of visual processing in age-related macular degeneration depends on foveal loss. *Optom Vis Sci*, 91, e199-206.
- Dragoi, V., Sharma, J., & Sur, M. (2000). Adaptation-induced plasticity of orientation tuning in adult visual cortex. *Neuron*, 28, 287-298.
- Garrett, A. T., Goosens, N. G., Rehner, N. J., Patterson, M. J., Harrison, J., Sammut, I., & Cotter, J. D. (2014). Short-term heat acclimation is effective and may be enhanced rather than impaired by dehydration. *Am J Hum Biol*, 26, 311-320.
- Gilbert, C. D., & Wiesel, T. N. (1992). Receptive field dynamics in adult primary visual cortex. *Nature*, 356, 150-152.
- Graham, N. V. S. (1989). *Visual Pattern Analyzers*. New York: Oxford University Press.
- Haak, K. V., Cornelissen, F. W., & Morland, A. B. (2012). Population receptive field dynamics in human visual cortex. *PLoS One*, 7, e37686.
- Haak, K. V., Fast, E., Baek, Y., & Mesik, J. (2014a). Equalization and decorrelation in primary visual cortex. *J Neurophysiol*, 112, 501-503.
- Haak, K. V., Fast, E., Bao, M., Lee, M., & Engel, S. A. (2014b). Four days of visual contrast deprivation reveals limits of neuronal adaptation. *Curr Biol*, 24, 2575-2579.
- Haak, K. V., Langers, D. R., Renken, R., van Dijk, P., Borgstein, J., & Cornelissen, F. W. (2014c). Abnormal visual field maps in human cortex: a mini-review and a case report. *Cortex*, 56, 14-25.
- Hirsch, H. V., & Spinelli, D. N. (1970). Visual experience modifies distribution of horizontally and vertically oriented receptive fields in cats. *Science*, 168, 869-871.
- Hochstein, S., & Ahissar, M. (2002). View from the top: hierarchies and reverse hierarchies in the visual system. *Neuron*, 36, 791-804.
- Huberman, A. D., & Niell, C. M. (2011). What can mice tell us about how vision works? *Trends Neurosci*, 34, 464-473.
- Kaas, J. H., Krubitzer, L. A., Chino, Y. M., Langston, A. L., Polley, E. H., & Blair, N. (1990). Reorganization of retinotopic cortical maps in adult mammals after lesions of the retina. *Science*, 248, 229-231.
- Keck, T., Mrsic-Flogel, T. D., Vaz Afonso, M., Eysel, U. T., Bonhoeffer, T., & Hubener, M. (2008). Massive restructuring of neuronal circuits during functional reorganization of adult visual cortex. *Nat Neurosci*, 11, 1162-1167.

- 1
2
3
4 Kohn, A. (2007). Visual adaptation: physiology, mechanisms, and functional benefits. *J*
5 *Neurophysiol*, 97, 3155-3164.
6
7
8 Komatsu, H. (2006). The neural mechanisms of perceptual filling-in. *Nat Rev Neurosci*, 7, 220-
9 231.
10
11 Maffei, L., Fiorentini, A., & Bisti, S. (1973). Neural correlate of perceptual adaptation to
12 gratings. *Science*, 182, 1036-1038.
13
14 Masuda, Y., Dumoulin, S. O., Nakadomari, S., & Wandell, B. A. (2008). V1 projection zone
15 signals in human macular degeneration depend on task, not stimulus. *Cereb Cortex*, 18,
16 2483-2493.
17
18
19 Masuda, Y., Horiguchi, H., Dumoulin, S. O., Furuta, A., Miyauchi, S., Nakadomari, S., &
20 Wandell, B. A. (2010). Task-dependent V1 responses in human retinitis pigmentosa. *Invest*
21 *Ophthalmol Vis Sci*, 51, 5356-5364.
22
23
24 Movshon, J. A., & Lennie, P. (1979). Pattern-selective adaptation in visual cortical neurones.
25 *Nature*, 278, 850-852.
26
27
28 Muckli, L., Naumer, M. J., & Singer, W. (2009). Bilateral visual field maps in a patient with
29 only one hemisphere. *Proc Natl Acad Sci U S A*, 106, 13034-13039.
30
31
32 Murakami, I., Komatsu, H., & Kinoshita, M. (1997). Perceptual filling-in at the scotoma
33 following a monocular retinal lesion in the monkey. *Vis Neurosci*, 14, 89-101.
34
35
36 Ohl, F. W., & Scheich, H. (2005). Learning-induced plasticity in animal and human auditory
37 cortex. *Curr Opin Neurobiol*, 15, 470-477.
38
39
40 Ohzawa, I., Sclar, G., & Freeman, R. D. (1985). Contrast gain control in the cat's visual system. *J*
41 *Neurophysiol*, 54, 651-667.
42
43
44 Patterson, C. A., Wissig, S. C., & Kohn, A. (2014). Adaptation disrupts motion integration in the
45 primate dorsal stream. *Neuron*, 81, 674-686.
46
47
48 Pettet, M. W., & Gilbert, C. D. (1992). Dynamic changes in receptive-field size in cat primary
49 visual cortex. *Proc Natl Acad Sci U S A*, 89, 8366-8370.
50
51
52 Polley, D. B., Steinberg, E. E., & Merzenich, M. M. (2006). Perceptual learning directs auditory
53 cortical map reorganization through top-down influences. *J Neurosci*, 26, 4970-4982.
54
55
56 Ramachandran, V. S., & Gregory, R. L. (1991). Perceptual filling in of artificially induced
57 scotomas in human vision. *Nature*, 350, 699-702.
58
59
60
61
62
63
64
65

- 1
2
3
4 Robertson, D., & Irvine, D. R. (1989). Plasticity of frequency organization in auditory cortex of
5 guinea pigs with partial unilateral deafness. *J Comp Neurol*, 282, 456-471.
6
7 Schmid, L. M., Rosa, M. G., & Calford, M. B. (1995). Retinal detachment induces massive
8 immediate reorganization in visual cortex. *Neuroreport*, 6, 1349-1353.
9
10 Schumacher, E. H., Jacko, J. A., Primo, S. A., Main, K. L., Moloney, K. P., Kinzel, E. N., &
11 Ginn, J. (2008). Reorganization of visual processing is related to eccentric viewing in
12 patients with macular degeneration. *Restor Neurol Neurosci*, 26, 391-402.
13
14 Schwartz, O., Hsu, A., & Dayan, P. (2007). Space and time in visual context. *Nat Rev Neurosci*,
15 8, 522-535.
16
17 Series, P., Stocker, A. A., & Simoncelli, E. P. (2009). Is the homunculus "aware" of sensory
18 adaptation? *Neural Comput*, 21, 3271-3304.
19
20 Smirnakis, S. M., Brewer, A. A., Schmid, M. C., Tolias, A. S., Schuz, A., Augath, M., Inhoffen,
21 W., Wandell, B. A., & Logothetis, N. K. (2005). Lack of long-term cortical reorganization
22 after macaque retinal lesions. *Nature*, 435, 300-307.
23
24 Sunness, J. S., Liu, T., & Yantis, S. (2004). Retinotopic mapping of the visual cortex using
25 functional magnetic resonance imaging in a patient with central scotomas from atrophic
26 macular degeneration. *Ophthalmology*, 111, 1595-1598.
27
28 Ullman, S., & Schechtman, G. (1982). Adaptation and gain normalization. *Proc R Soc Lond B*
29 *Biol Sci*, 216, 299-313.
30
31 Wainwright, M. J. (1999). Visual adaptation as optimal information transmission. *Vision Res*, 39,
32 3960-3974.
33
34 Wandell, B. A., & Smirnakis, S. M. (2009). Plasticity and stability of visual field maps in adult
35 primary visual cortex. *Nat Rev Neurosci*, 10, 873-884.
36
37 Wang, Q., & Burkhalter, A. (2007). Area map of mouse visual cortex. *J Comp Neurol*, 502, 339-
38 357.
39
40 Weil, R. S., & Rees, G. (2011). A new taxonomy for perceptual filling-in. *Brain Res Rev*, 67, 40-
41 55.
42
43 Werth, R. (2006). Visual functions without the occipital lobe or after cerebral hemispherectomy
44 in infancy. *Eur J Neurosci*, 24, 2932-2944.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65